

ORIGINAL ARTICLE

Antioxidant Effect of Extra Virgin Olive Oil and Apple Cider Vinegar in Type 2 Diabetic Rat ModelRuqaiyya Nazir^{1*}, Muniza Saeed², Hifza Noor Lodhi^{3*}, Ghulam Mustafa⁴, Seher Naeem⁵, Zunnera Rashid Chaudhry⁶**ABSTRACT**

Objective: Comparison between the antioxidant effects of extra virgin olive oil and apple cider vinegar in rats induced with diabetes mellitus type 2.

Study Design: Randomized controlled trial.

Place and Duration of Study: This Study was conducted in Postgraduate Medical Institute, Animal House Lahore, Pakistan from May 2021 to June 2021.

Methods: In this study, 40 male Sprague Dawley rats were divided into 4 groups i.e., Group I was NC (negative control), Group II PC (positive control), Group III EVOO (Extra virgin olive oil) and Group IV (Apple cider vinegar), each group having 10 rats. Diabetes was induced in all rats except the rats of NC group at the start of the study by intraperitoneal administration of injection nicotinamide, followed by injection Streptozosin (STZ) after 15 minutes. Group III was given 1ml/100gBW/ day EVOO and Group IV was given 2ml/kg BW/day diluted ACV with distilled water in 1:5 orally for 4 weeks. Sampling was done after 4 weeks for determination of oxidative stress markers Malonaldehyde, Superoxide dismutase and Total antioxidant status in serum.

Results: Intake of EVOO and ACV showed reduction in serum Malonaldehyde levels as compared to positive control group with *P*-values 0.000 and 0.014 respectively. Serum superoxide dismutase activity and Total antioxidant status were lowest in the positive control group while both the treatment groups showed significant enhancement in these parameters as compared to positive control group with *P*-value = 0.000.

Conclusion: Both extra virgin olive oil and apple cider vinegar have antioxidant effects in type 2 diabetic rats. However, Extra virgin olive oil is more potent.

Keywords: Antioxidant Effects, Diabetes Mellitus, Olive Oil.

How to cite this: Nazir R, Saeed M, Lodhi HN, Mustafa G, Naeem S, Chaudry ZR. Antioxidant Effect of Extra Virgin Olive Oil and Apple Cider Vinegar in Type 2 Diabetic Rat Model. *Life and Science*. 2024; 5(4): 557-562. doi: <http://doi.org/10.37185/LnS.1.1.780>

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International license. (<https://creativecommons.org/licenses/by-nc/4.0/>). Non-commercial uses of the work are permitted, provided the original work is properly cited.

¹Department of Physiology
Islamic International Medical Collage Rawalpindi, Pakistan

²Department of Physiology
Postgraduate Medical Institute Lahore, Pakistan

³Department of Physiology
Rashid Latif Khan University Lahore, Pakistan

⁴Department of Nephrology
Armed Forces Institute of Urology (AFIU) Rawalpindi
Pakistan

⁵Department of Physiology
Combined Military Hospital Rawalpindi (CMH) Lahore
Pakistan

⁶Department of Pharmacology
Akhtar Saeed Medical & Dental College Rawalpindi
Pakistan

Correspondence:

Dr. Hifza Noor Lodhi
Associate Professor, Physiology
Rashid Latif Khan University Lahore, Pakistan
E-mail: noor.lodhi@rlkumc.edu.pk

Received: Jun 30, 2024; Revised: Aug 30, 2024
Accepted: Sep 15, 2024

Introduction

Type 2 diabetes mellitus is differentiated by resistance to insulin at the insulin receptors and changed metabolism and dyslipidemia.¹ Diabetes mellitus has surfaced as an issue since past few eras, leading to colossal drain on health resources.² There were estimated 451 million people diabetics around the world and 693 million to have it by the 2045.³ An estimated 26.3% adults are suffering from this condition in Pakistan.⁴

In diabetes, persistent high blood sugar levels leading to the generation of ROS. The imbalance between free radicals and antioxidants in individuals with diabetes contributes to the harm of various organs.⁵ Free radicals cause alterations in structure and functions of cellular components i.e., cell

membrane and biomolecules such as lipids, nucleic acid and proteins. Malondialdehyde (MDA) is produced by lipid peroxidation of membrane lipids and is a prime marker of cell injury. Lipid peroxidation is involved in development of micro and macro vascular complications of diabetes. Oxidative damage is prevented by enzymes such as catalase (CAT), glutathione peroxidase and superoxide dismutase (SOD) and non-enzymatic components of total antioxidant status (TAS) for example ascorbic acid and tocopherol but there is a significant decline TAS in DM.⁶

Diabetic patients are currently being treated by insulin and oral hypoglycemic agents. The challenges associated with insulin use and the side effects of medications are driving an increased interest in natural treatment options for diabetes.⁷ Among these are phenolic compounds, which are recognized for their hypoglycemic and antioxidant properties. Extra virgin olive oil is particularly high in phenolic compounds compared to other types of olive oil.⁸ Apple cider vinegar, which is made through a two-step fermentation process, contains 5-6% acetic acid.⁹ The phenolic compounds in ACV inhibit generation of ROS antioxidant and lower blood glucose levels.¹⁰ Additionally, apple cider vinegar exhibits hepatoprotective properties due to phloretin, which plays a crucial role in non-enzymatic protection against oxidative stress-induced liver damage.¹¹

This study is aimed to compare the antioxidant effect of EVOO and ACV in diabetic rats by determining serum levels of oxidative stress markers.

Methods

This experimental study, with a duration of 40 days, was conducted at the Animal House of Postgraduate Medical Institute, Lahore from May 2021 to June 2021. Approval was taken by Ethical Committee of the institute on dated: 14th March 2021 vide letter no: 00/20/5/2019.

Sample size was calculated by using formula;

$$n = \frac{\sigma^2(z_{1-\alpha/2} + z_{1-\beta})^2}{(\mu_o - \mu_a)^2}$$

The level of significance (α) = 5% and level of confidence = 95%. Values of μ_o & μ_a were obtained from ALT levels from the result of Balamash KS et al. Sample size calculated was 10 in each group.¹²

40 adult healthy male rats of (180-200 gram) selected by simple random sampling were bought from local market in Lahore. Rats with any ailment signs were excluded from the study.

Before starting the experiments, the rats were allowed a week to acclimate in the animal house. They were in appropriately sized, cages at a room temperature of 22-24°C and provided with rat chow and water orally. Following this acclimatization period, the rats were randomly assigned into four groups (10 rats per group).

Type 2 diabetes mellitus was induced in all rats except those in the normal control by administering intraperitoneal injections of nicotinamide using a 26-gauge needle, followed by an injection of streptozotocin (STZ). Rats with blood glucose levels exceeding 200 mg/dL, as measured by a handheld glucometer, 72 hours after the STZ injection, were considered diabetic.¹³

Both control groups, Group I (NC) and Group II (PC), received 1 ml/100 g body weight/day of distilled water orally. Group III (EVOO) was administered 1 ml/100 g body weight/day of extra virgin olive oil (Spain), while Group IV (ACV) was given 2 ml/kg body weight/day of diluted apple cider vinegar mixed with distilled water.¹²⁻¹⁴

At the end of 4th week cardia puncture was performed to collect 4 ml blood of each rat after overnight fast. Light anesthesia with chloroform was used to collect samples by using 26-gauge needle. Sample tubes were placed in straight standing position for 30 minutes at room temperature and then centrifuged. Serum MDA levels were measured by ELISA while serum SOD & TAS were measured by colorimetric method.

Results

Very highly significant difference in serum stress markers (MDA, SOD and TAS) levels was noted when comparison among groups was done by Kruskal Wallis *H* test ($P=0.000$) (Table-1) because data was non normally distributed. Serum MDA levels of NC rats were significantly less than the PC rats with a $P=0.000$. Serum MDA levels of ACV and EVOO groups were significantly less than PC group with P -values of 0.000 and 0.014 respectively. (Table-2).

Serum SOD levels of NC rats were higher than the PC rats with $P=0.000$. Serum TAS of PC rats were very

highly significantly lower than the NC rats with a $P = 0.000$. Comparison of serum Total Antioxidant Status

of both the groups with Positive Control group had P -value of 0.0000 . (Table- 2).

Table-1: Comparison of serum stress markers (malondialdehyde, superoxide dismutase and total antioxidant status) levels

Groups	Malondialdehyde (nmol/mL)	Superoxide dismutase (U/mL)	Total antioxidant status (U/mL)
Median	1.35	22.80	4.25
(IQR)	(1.20-1.52)	(22.80-23.62)	(3.92-4.47)
Median	4.30	11.70	1.24
(IQR)	(3.87-4.82)	(11.20-12.65)	(0.49-2.10)
Median	2.50	17.20	4.68
(IQR)	(2.20-2.57)	(16.67-17.77)	(4.53-6.41)
Median	3.50	16.60	3.82
(IQR)	(3.27-3.92)	(16.15-17.05)	(2.99-4.07)
P -value	0.000***	0.000***	0.000***

*** P value very highly significant. PC: Positive control NC: Normal control, EVOO: Extra virgin Olive oil, ACV: Apple Cider vinegar

Table-2: Multiple comparison of serum stress markers (malondialdehyde, superoxide dismutase and total antioxidant status) levels among groups

Groups	Malondialdehyde (nmol/mL)	Superoxide dismutase (U/mL)	Total antioxidant status (U/mL)
Group I (NC)	Median 1.35 (IQR) (1.20-1.52) $^2p=0.000$ ***	Median 22.80 (IQR) (22.80-23.62) $^2p=0.000$ ***	Median 4.25 (IQR) (3.92-4.47) $^2p=0.000$ ***
Group II (PC)	Median 4.30 (IQR) (3.87-4.82)	Median 11.70 (IQR) (11.20-12.65)	Median 1.24 (IQR) (0.49-2.10)
Group III (EVOO)	Median 2.50 (IQR) (2.20-2.57) $^1p=0.000$ *** $^2p=0.000$ ***	Median 17.20 (IQR) (16.67-17.77) $^1p=0.000$ *** $^2p=0.000$ ***	Median 4.68 (IQR) (4.53-6.41) $^1p=0.006$ ** $^2p=0.000$ ***
Group IV (ACV)	Median 3.50 (IQR) (3.27-3.92) $^1p=0.000$ *** $^2p=0.014$ *	Median 16.60 (IQR) (16.15-17.05) $^1p=0.000$ *** $^2p=0.000$ ***	Median 3.82 (IQR) (2.99-4.07) $^1p=0.028$ * $^2p=0.000$ ***

* P -value significant. PC: Positive control NC: Normal control, EVOO: Extra virgin Olive oil, ACV: Apple Cider vinegar

Discussion

The Oxidative stress has a crucial part in pathogenesis of T2DM and its complications. This study was conducted to study the antioxidant effect of ACV and EVOO in rats induced with diabetes. There are many markers of oxidative stress including ROS themselves. Since ROS have short half-lives, it is appropriate to determine oxidative stress by measuring the products of oxidation reactions. Free radicals react with lipid bilayer of plasma membrane resulting in lipid peroxidation and MDA is produced as a result which is more stable than ROS.^{15,16}

In our study, DC group showed the highest levels of serum MDA among all the groups. Both the groups that were given treatment i.e., EVOO and ACV showed reduction in serum MDA levels as compared to DC group with p values of 0.000 and 0.014 respectively. Serum SOD activity and serum TAS were lowest in the DC group among all the groups. While both the groups that were treated had very highly significant enhancement in serum SOD activity as well as TAS (P values < 0.001) as compared to DC group. Significantly raised MDA levels were noted in T2DM

with and without complications in previous studies, confirming results of our study.^{15,16} Diabetes lowered the activities of antioxidant enzymes, such as activity of SOD in the study of Karabag-Coban F et al. It was suggested that reduction in activity of enzymes is due to their enhanced utilization in combating oxidative stress in diabetic group. Another study is in accordance with our study, reporting decrease in plasma SOD activity as well as TAS in type 2 diabetics as compared to normal participants.¹⁷

It was suggested that reduction in activity of enzymes is due to their enhanced utilization in combating oxidative stress in diabetic group. Another study is in accordance with our study, reporting decrease in plasma SOD activity as well as TAS in type 2 diabetics as compared to normal participants.¹⁸

Olive oil contains abundant bio phenols like HT, oleuropein and their derivatives which have antioxidative effect.¹⁹ Olive oil helps to increase serum total antioxidant capacity by enhancing the level and activity of antioxidant enzymes. Olive oil consumption leads to increase in MUFA content in LDL, MUFA are more resistant to oxidation than polyunsaturated fatty acids, preserving antioxidant compounds bound to LDL.²⁰ Temiz and Temur reported that polyphenolic compounds present in olive leaf extract increased the activity of antioxidant enzymes like SOD, CAT and glutathione activity in the diabetic rats of their study. It was suggested that these polyphenols also increase the expression of enzymes SOD and CAT at the level of transcription.²¹ Karabag-Coban F et al. indicated decreased lipid peroxidation resulting in decreased MDA levels in oleuropein treated diabetic animals. They attributed this achievement to the ability of oleuropein (an important component of olive oil) to decompose and trap free radicals before reaching the cellular targets. Moreover, oleuropein was found to reestablish SOD activity in the treatment group of their study.¹⁷ However, Atefi M et al. noted non-significant effect of olive oil on MDA level in type 2 diabetics; they have attributed their results to short duration of their study.²²

The antioxidant effects of ACV shown in our study are in accordance to the properties of ACV reported by previous studies. Polyphenols present in vinegars

provide antioxidant properties to vinegars because they have capacity to remove free radicals, reduce oxidants and chelate transition metal ions. The benzene ring of polyphenols blocks free radical chains and the hydroxyl group on benzene ring chelate transition metal ions, preventing oxidation reaction. Moreover, phenolic hydroxyl group can be oxidized to quinone, thus polyphenols also have a reducing effect.²³ Gheflati A et al. noted in their study that consumption of apple vinegar by type 2 diabetics had inverse relation with MDA level and they attributed this finding to the ability of flavonoids to scavenge ROS. Moreover, total antioxidant capacity of the vinegar group was also assessed in their study which was enhanced as compared to diabetic control group, confirming our results.²⁴ Abdulrauf RA et al. observed and reported in their study that decline in lipid peroxidation as evidenced by fall in serum MDA level by treating oxidative stress by ACV and they suggested catechins present in ACV to be responsible for reduction in LDL oxidation. Their treatment group also showed an increase in serum SOD activity, attributed to up regulation of the release of endogenous SOD by ACV as well as to the enhancement of SOD synthesis by the minerals (potassium, zinc and copper) present in ACV. Moreover, it was also suggested that phenolic acids and vitamins in ACV can remove superoxide anion, resulting in potent antioxidant activity.²⁵

Conclusion

This study concludes that apple cider vinegar as well as extra virgin olive oil have antioxidant effect in rat model of type because both lowered the oxidative stress marker and increasing the antioxidants. The effect of EVOO was more definite.

Limitations: Large clinical trials are needed to confirm these findings in human settings.

Acknowledgment: We did not use Artificial intelligence assisted technologies in the creation of this work.

Conflict of Interest: The authors declare no conflict of interest.

Grant Support and Financial Disclosure: None.

REFERENCES

1. Rom S, Zuluaga-Ramirez V, Gajghate S, Seliga A, Winfield M,

- Heldt NA et al. Hyperglycemia-driven neuroinflammation compromises BBB leading to memory loss in both diabetes mellitus (DM) type 1 and type 2 mouse models. *Molecular Neurobiology*. 2019; 56: 1883-96. doi: 10.1007/s12035-018-1195-5
2. Kausar S, Humayun A, Ahmed Z, Abbas MA, Tahir A. Effect of apple cider vinegar on glycemic control, hyperlipidemia and control on body weight in type 2 diabetes patients. *International Journal of Medical Research and Health Sciences*. 2019; 8: 59-74.
 3. Cho NH, Shaw JE, Karuranga S, Huang Y, da Rocha Fernandes JD, Ohlrogge AW, et al. IDF Diabetes Atlas: Global estimates of diabetes prevalence for 2017 and projections for 2045. *Diabetes Research and Clinical Practice*. 2018; 138: 271-81. doi: 10.1016/j.diabres.2018.02.023
 4. Ijaz M, Ali I, Hussain A. Diabetes mellitus in Pakistan: the past, present, and future. *International Journal of Diabetes in Developing Countries*. 2020; 40: 153-4. doi: 10.1007/s13410-019-00754-x1
 5. Dal S, Sigrist S. The protective effect of antioxidants consumption on diabetes and vascular complications. *Diseases*. 2016; 4: 24. doi: 10.3390/diseases4030024
 6. Rehman K, Akash MS. Mechanism of generation of oxidative stress and pathophysiology of type 2 diabetes mellitus: how are they interlinked? *Journal of cellular biochemistry*. 2017; 118: 3577-85. doi: 10.1002/jcb.26097
 7. Laaboudi WA, Ghanam JA, Ghomari OU, Sounni FA, Merzouki MO, Benlemlih MO. Hypoglycemic and hypolipidemic effects of phenolic olive tree extract in streptozotocin diabetic rats. *International Journal of Pharmacy and Pharmaceutical Sciences*. 2016; 8: 287-91. doi: 10.22159/ijpps.2016v8i12.14077
 8. Saibandith B, Spencer JP, Rowland IR, Commane DM. Olive polyphenols and the metabolic syndrome. *Molecules*. 2017; 22: 1082. doi: 10.3390/molecules22071082
 9. Fathy SM, Drees EA. Protective effects of Egyptian cloudy apple juice and apple peel extract on lipid peroxidation, antioxidant enzymes and inflammatory status in diabetic rat pancreas. *BMC complementary and alternative medicine*. 2016; 16: 8. doi:10.1186/s12906-015-0957-0
 10. Morgan J, Mosawy S. The potential of apple cider vinegar in the management of type 2 diabetes. *International Journal of Diabetes Research*. 2016; 5: 129-34. doi: 10.5923/j.diabetes.20160506.02
 11. Lu Y, Chen J, Ren D, Yang X, Zhao Y. Hepatoprotective effects of phloretin against CCl₄-induced liver injury in mice. *Food and Agricultural Immunology*. 2017; 28: 211-22. doi: 10.1080/09540105.2016.1258546
 12. Balamash KS, Alkreathy HM, Al Gahdali EH, Khoja SO, Ahmad A. Comparative biochemical and histopathological studies on the efficacy of metformin and virgin olive oil against streptozotocin-induced diabetes in Sprague-Dawley rats. *Journal of Diabetes Research*. 2018; 2018: 4692197. doi: 10.1155/2018/4692197
 13. Cam ME, Hazar-Yavuz AN, Yildiz S, Ertas B, Adakul BA, Taskin T et al. The methanolic extract of *Thymus praecox* subsp. *skorpilii* var. *skorpilii* restores glucose homeostasis, ameliorates insulin resistance and improves pancreatic β -cell function on streptozotocin/nicotinamide-induced type 2 diabetic rats. *Journal of ethnopharmacology*. 2019; 231: 29-38. doi: 10.1016/j.jep.2018.10.028
 14. Omar NA, Allithy AN, El Sayed SM. Hepatoprotective and antidiabetic effects of apple cider vinegar (a prophetic medicine remedy) on the liver of male rats. *The Egyptian Journal of Hospital Medicine*. 2016; 62: 95-104. doi: 10.12816/0021418
 15. Mishra S, Mishra BB. Study of lipid peroxidation, nitric oxide end product, and trace element status in type 2 diabetes mellitus with and without complications. *International Journal of Applied and Basic Medical Research*. 2017; 7: 88-93. doi: 10.4103/2229-516X.205813
 16. Bigagli E, Lodovici M. Circulating oxidative stress biomarkers in clinical studies on type 2 diabetes and its complications. *Oxidative medicine and cellular longevity*. 2019; 2019: 5953685. doi: 10.1155/2019/5953685
 17. Karabag-Coban F, Hazman O, Bozkurt M, Ince S. Antioxidant status and anti-inflammatory effects of oleuropein in streptozotocin-induced diabetic nephropathy in rats. *European Journal of Medicinal Plants*. 2017; 18: 1-10. doi: 10.9734/EJMP/2017/31953
 18. Mohammed AK, Khurshid CA, Abdulrahman HB. Uncompensated response to the oxidation stress of the diabetes in both male and female patients with type II diabetes. *Eurasian Journal of Biosciences*. 2020; 14: 3139-44.
 19. Alkhatib A, Tsang C, Tuomilehto J. Olive oil nutraceuticals in the prevention and management of diabetes: From molecules to lifestyle. *International Journal of Molecular Sciences*, 2018; 19: 2024. doi: 10.3390/ijms19072024
 20. Ortner Hadžiabdić M, Vitali Čepo D, Rahelić D, Božikov V. The effect of the Mediterranean diet on serum total antioxidant capacity in obese patients: a randomized controlled trial. *Journal of the American College of Nutrition*, 2016; 35: 224-35. doi: 10.1080/07315724.2014.982770

21. Temiz MA, Temur A. Olive (*Olea europaea* L.) leaf extract boosts antioxidant status and attenuates hepatic and pancreatic changes in diabetic rats. *Applied Biological Research*. 2019; 21: 10-9. doi: 10.5958/0974-4517.2019.00002.8
22. Atefi M, Pishdad GR, Faghih S. The effects of canola and olive oils on insulin resistance, inflammation and oxidative stress in women with type 2 diabetes: a randomized and controlled trial. *Journal of Diabetes & Metabolic Disorders*, 2018; 17: 85-91. doi: 10.1007/s40200-018-0343-9
23. Chen H, Chen T, Giudici P, Chen F. Vinegar functions on health: Constituents, sources, and formation mechanisms. *Comprehensive Reviews in Food Science and Food Safety*, 2016; 15: 1124-38. doi: 10.1111/1541-4337.12228
24. Gheflati A, Bashiri R, Ghadiri-Anari A, Reza JZ, Kord MT, Nadjarzadeh A. The effect of apple vinegar consumption on glycemic indices, blood pressure, oxidative stress, and homocysteine in patients with type 2 diabetes and dyslipidemia: A randomized controlled clinical trial. *Clinical Nutrition*, 2019; 33: 132-8. doi: 10.1016/j.clnesp.2019.06.006
25. Abdulrauf RA, Dawud FA, Emmanuel NS, Muhammad HD, Dange AS, David BA, et al. Lipid peroxidation and some antioxidant enzymes evaluation in apple cider vinegar (ACV) treated male and female Wistar rats exposed to chronic restraint stress. *Advances in Enzyme Research*. 2018; 6: 21-8. doi: 10.4236/aer.2018.63003

Authors Contribution

RN: Idea conception, study designing, data collection, data analysis, results and interpretation

MS: Idea conception, study designing, data collection, data analysis, results and interpretation

HNL: Idea conception, study designing, data collection, manuscript writing and proofreading

GM: Idea conception, study designing, data collection

SN: Idea conception, study designing, data collection

ZRC: Data analysis, results and interpretation, manuscript writing and proofreading

.....